

## Consanguinity, Founder Effect

- Abney, M., McPeck, M. S. and Ober, C. (2001). Broad and narrow heritabilities of quantitative traits in a founder population. *American Journal of Human Genetics* **68**: 1302-1307.
- Adato, A., Raskin, L., Petit, C. and Bonne-Tamir, B. (2000). Deafness heterogeneity in a Druze isolate from the Middle East: novel OTOF and PDS mutations, low prevalence of GJB2 35delG mutation and indication for a new DFNB locus. *European Journal of Human Genetics* **8**: 437-442.
- Agarwala, R., Schaffer, A. A. and Tomlin, J. F. (2001). Towards a complete North American Anabaptist Genealogy II: analysis of inbreeding. *Human Biology* **73**: 533-545.
- Alos, N., Moisan, A. M., Ward, L., Desrochers, M., Legault, L., Leboeuf, G., Van Vliet, G. and Simard, J. (2000). A novel A10E homozygous mutation in the HSD3B2 gene causing severe salt-wasting 3beta-hydroxysteroid dehydrogenase deficiency in 46,XX and 46,XY French-Canadians: evaluation of gonadal function after puberty. *Journal of Clinical Endocrinology and Metabolism* **85**: 1968-1974.
- Baumann, P., Myllylä, V. V. and Leisti, J. (1998). Myotonia congenita in northern Finland: an epidemiological and genetic study. *Journal of Medical Genetics* **35**: 293-296.
- Bejjani, B. A., Lewis, R. A., Tomey, K. F., Anderson, K. L., Dueker, D. K., Jabak, M., Astle, W. F., Otterud, B., *et al.* (1998). Mutations in CYP1B1, the gene for cytochrome P4501B1, are the predominant cause of primary congenital glaucoma in Saudi Arabia. *American Journal of Human Genetics* **62**: 325-333.
- Ben Arab, S., Hmani, M., Denoyelle, F., Boulila-Elgaied, A., Chardenoux, S., Hachicha, S., Petit, C. and Ayadi, H. (2000). Mutations of GJB2 in three geographic isolates from northern Tunisia: evidence for genetic heterogeneity within isolates. *Clinical Genetics* **57**: 439-443.
- Brunkow, M. E., Gardner, J. C., Van Ness, J., Paeper, B. W., Kovacevich, B. R., Proll, S., Skonier, J. E., Zhao, L., *et al.* (2001). Bone dysplasia sclerosteosis results from loss of the SOST gene product, a novel cystine knot-containing protein. *American Journal of Human Genetics* **68**: 577-589.
- Bull, L. N., Roche, E., Song, E. J., Pedersen, J., Knisely, A. S., van Der Hagen, C. B., Eiklid, K., Aagenaes, O., *et al.* (2000). Mapping of the locus for cholestasis-lymphedema syndrome (Aagenaes syndrome) to a 6.6-cM interval on chromosome 15q. *American Journal of Human Genetics* **67**: 994-999.
- De Braekeleer, M. and Dao, T. (1994). Hereditary disorders in the French Canadian population of Quebec. II. Contribution of Perche. *Human Biology* **66**: 225-249.

- De Braekeleer, M. and Dao, T. (1994). Hereditary disorders in the French Canadian population of Quebec. I In search of founders. *Human Biology* **66**: 205-223.
- Ding, C., Buckingham, B. and Levine, M. A. (2001). Familial isolated hypoparathyroidism caused by a mutation in the gene for the transcription factor GCMB. *Journal of Clinical Investigation* **108**: 1215-1220.
- Hart, T. C., Hart, P. S., Michalec, M. D., Zhang, Y., Firatli, E., Van Dyke, T. E., Stabholz, A., Zlotogorski, A., *et al.* (2000). Haim-Munk syndrome and Papillon-Lefevre syndrome are allelic mutations in cathepsin C. *Journal of Medical Genetics* **37**: 88-94.
- Ianakiev, P., Kilpatrick, M. W., Daly, M. J., Zolindaki, A., Bagley, D., Beighton, G., Beighton, P. and Tsiouras, P. (2000). Localization of an acromesomelic dysplasia on chromosome 9 by homozygosity mapping. *Clinical Genetics* **57**: 278-283.
- Moulard, B., Genton, P., Grid, D., Jeanpierre, M., Ouazzani, R., Mrabet, A., Morris, M., LeGuern, E., *et al.* (2002). Haplotype study of West European and North African Unverricht-Lundborg chromosomes: evidence for a few founder mutations. *Human Genetics* **111**: 255-262.
- O'Brien, E., Kerber, R. A., Jorde, L. B. and Rogers, A. R. (1994). Founder effect: assessment of variation in genetic contributions among founders. *Human Biology* **66**: 185-204.
- Papponen, H., Toppinen, T., Baumann, P., Myllylä, V., Leisti, J., Kuivaniemi, H., Tromp, G. and Myllylä, R. (1999). Founder mutations and the high prevalence of myotonia congenita in northern Finland. *Neurology* **53**: 297-302.
- Peltonen, L. (1997). Molecular background of the Finnish disease heritage. *Annals of Medicine* **29**: 553-556.
- Peltonen, L., Jalanko, A. and Varilo, T. (1999). Molecular genetics of the Finnish disease heritage. *Human Molecular Genetics* **8**: 1913-1923.
- Picard, C., Fieschi, C., Altare, F., Al-Jumaah, S., Al-Hajjar, S., Feinberg, J., Dupuis, S., Soudais, C., *et al.* (2002). Inherited interleukin-12 deficiency: IL12B genotype and clinical phenotype of 13 patients from six kindreds. *American Journal of Human Genetics* **70**: 336-348.
- Rudan, I., Ranzani, G. N., Strnad, M., Vorko-Jovic, A., John, V., Unusic, J., Kern, J., Ivankovic, D., *et al.* (1999). Surname as 'cancer risk' in extreme isolates: example from the island of Lastovo, Croatia. *Collegium Antropologicum* **23**: 557-569.
- Scriver, C. R. (2001). Human genetics: lessons from Quebec populations. *Annual Review of Genomics and Human Genetics* **2**: 69-101.

- Tullio-Pelet, A., Salomon, R., Hadj-Rabia, S., Mugnier, C., de Laet, M. H., Chaouachi, B., Bakiri, F., Brottier, P., *et al.* (2000). Mutant WD-repeat protein in triple-A syndrome. *Nature Genetics* **26**: 332-335.
- Virtaneva, K. (1996). Progressive myoclonus epilepsy EPM1 locus maps to a 175-kb interval in distal 21q. *American Journal of Human Genetics* **58**: 1247-1253.
- Visapaa, I., Salonen, R., Varilo, T., Paavola, P. and Peltonen, L. (1999). Assignment of the locus for hydrolethalus syndrome to a highly restricted region on 11q23-25. *American Journal of Human Genetics* **65**: 1086-1095.
- Wiszniewski, W., Fondaneche, M. C., Lambert, N., Masternak, K., Picard, C., Notarangelo, L., Schwartz, K., Bal, J., *et al.* (2000). Founder effect for a 26-bp deletion in the RFXANK gene in North African major histocompatibility complex class II-deficient patients belonging to complementation group B. *Immunogenetics* **51**: 261-267.
- Young, T. L., Woods, M. O., Parfrey, P. S., Green, J. S., Hefferton, D. and Davidson, W. S. (1999). A founder effect in the Newfoundland population reduces the Bardet-Biedl syndrome I (BBS1) interval to 1 cM. *American Journal of Human Genetics* **65**: 1680-1687.